

# Infant formula – An overview

August 2017

## **Infant formula – An overview.** August 2017

Dr Helen Crawley and Susan Westland

© First Steps Nutrition Trust, 2017

### **DISCLAIMER**

The opinions given in this resource are those of the authors. We have reviewed the evidence provided by manufacturers, and the claims made on their websites, and have summarised our thoughts about this data and how these claims match current expert opinion from respected scientific bodies. We strongly recommend that health professionals review the evidence for themselves and make their own minds up about whether the statements and claims made by companies about their products are scientific and factual.

This report is provided for information only and individual advice on diet and health should always be sought from appropriate health professionals.

## **First Steps Nutrition Trust**

First Steps Nutrition Trust is a charity which provides clear, evidence-based and independent information and support for good nutrition from pre-conception to five years of age.

### **First Steps Nutrition Trust**

112 Queens Road

London SW19 8LS

[www.firststepsnutrition.org](http://www.firststepsnutrition.org)

E: [Helen@firststepsnutrition.org](mailto:Helen@firststepsnutrition.org)

Registered charity number: 1146408

# Contents

---

Introduction	5
Aptamil 1 First Milk	10
Aptamil Profutura 1 First Infant Milk	15
Bebivita First Infant Milk	21
Castlemil First Infant Milk	22
Cow & Gate 1 First Infant Milk	23
HiPP Organic Combiotic First Infant Milk	29
Holle Organic Infant Formula 1	34
Holle Organic Infant Goat Milk Formula 1	35
Kabrita Gold 1	36
Kendamil 1 First Infant Milk	38
Kendamil Mehadrin First Infant Milk	39
Mamia First Infant Milk	40
NANNYcare First Infant Milk	41
Similac First Infant Milk	43
SMA Pro First Infant Milk	44
SMA Wysoy	49
Hungry baby milks	50
Appendix Cost of infant formula in the UK July 2017	52

## List of Tables

TABLE 1	Macro and micronutrient requirements for infant formula and follow-on formula	6
TABLE 2	Key points made by EFSA about infant formula	7
TABLE 3	Infant milk products discussed in this resource	9
TABLE 4	Key components of Aptamil 1 First Milk	10
TABLE 5	Key components of Aptamil Profutura 1 First Infant Milk	15
TABLE 6	Key components of Bebivita First Infant Milk	21
TABLE 7	Key components of Castlemil First Infant Milk	22
TABLE 8	Key components of Cow & Gate 1 First Infant Milk	23
TABLE 9	Key components of HiPP Organic Combiotic First Infant Milk	29
TABLE 10	Key components of Holle Organic Infant Formula 1	34
TABLE 11	Key components of Holle Organic Infant Goat Milk Formula 1	35
TABLE 12	Key components of Kabrita Gold 1	36
TABLE 13	Key components of Kendamil 1 First Infant Milk	38
TABLE 14	Key components of Kendamil Mehadrin First Infant Milk	39
TABLE 15	Key components of Mamia First Infant Milk	40
TABLE 16	Key components of NANNYcare First Infant Milk	41
TABLE 17	Key components of Similac First Infant Milk	43
TABLE 18	Key components of SMA PRO First Infant Milk	44
TABLE 19	Key components of SMA Wysoy	49
TABLE 20	Summary of composition of hungry baby milks (powdered formulations only)	51
TABLE 21	Cost of infant formula in the UK – July 2017	52

# Introduction

---

## What information is in this resource?

This resource provides information about **infant formula**. These are breastmilk substitutes which have to meet the current UK Infant Formula Regulations, and which can be used from birth as the sole source of nutrition. In the UK we recommend these are used throughout the first year of life if infants are not breastfed, receiving breastmilk or if they are partially breastfed.

## What is the aim of this resource?

Despite regulations which state that all infant formula marketed in the UK must be of an agreed nutritional composition, and be appropriately labelled, products within the market may appear to be quite different. Manufacturers attempt to gain brand loyalty for their product through making claims about superiority of ingredients. Company representatives, mailings, websites and conference exhibitors can provide confusing information for health professionals, and the evidence the manufacturers present may appear convincing, and contradictory.

As health professionals may be asked about how infant formula differ, this resource provides an independent assessment of the products currently available in the UK.

This resource does not provide information about follow-on formula (which are not needed) or specialist formula. Information about all infant milks marketed in the UK, current regulations and an assessment of infant formula composition can be found in the report *Infant Milks in the UK* at [www.firststepsnutrition.org](http://www.firststepsnutrition.org).

## A brief summary of current regulations and expert views on composition

The compositional requirements are currently determined by the Infant Formula and Follow-on Formula Regulations (2007) and any amendments to those regulations. New regulations in the EU as part of delegated acts to the Foods for Special Groups Directive (EC 609/2013) have been published, but manufacturers have until 2020 to bring products in line with the new requirements and we will report on how this might affect products in future reports. The current compositional requirements which infant formula have been based on since 2007 are shown in Table 1. These have been recalculated to show them per 100ml of formula.

**TABLE 1**  
**Macro and micronutrient requirements for infant formula and follow-on formula**

<b>Regulations for infant formula (re-calculated by volume)</b>		
<b>MACRONUTRIENTS</b>	<b>Min/100ml</b>	<b>Max/100ml</b>
<b>Energy kJ</b>	250	295
kcal	60	70
<b>Protein g</b>	1.2	2.0
<b>Carbohydrate g</b>	6.0	9.4
of which lactose g	3.0	N/S
<b>Fat g</b>	2.9	4.0
<b>Linoleic acid mg</b>	200	800
<b>Linolenic acid mg</b>	33.5	N/S
<b>VITAMINS</b>		
<b>Vitamin A µg-RE</b>	40	121
<b>Vitamin C mg</b>	10	20.1
<b>Vitamin E mg</b>	0.34*	3.35
<b>Vitamin D µg</b>	0.67	1.68
<b>Vitamin K µg</b>	2.68	16.75
<b>Thiamin (B<sub>1</sub>) µg</b>	40.2	201
<b>Riboflavin (B<sub>2</sub>) µg</b>	53.6	268
<b>Niacin µg</b>	201	1005
<b>Vitamin B<sub>6</sub> µg</b>	23.45	117.25
<b>Vitamin B<sub>12</sub> µg</b>	0.07	0.34
<b>Folic acid µg</b>	6.7	33.5
<b>Biotin µg</b>	1.01	5.03
<b>Pantothenic acid µg</b>	268	1340
<b>MINERALS</b>		
<b>Calcium mg</b>	34	94
<b>Chloride mg</b>	33.5	107.2
<b>Copper µg</b>	23.45	67
<b>Fluoride µg</b>	N/S	67
<b>Iodine µg</b>	6.7	33.5
<b>Iron** mg</b>	0.2	0.87
<b>Magnesium mg</b>	3.35	10
<b>Manganese µg</b>	0.67	67
<b>Phosphorus**mg</b>	16.8	60.3
<b>Potassium mg</b>	40.2	107.2
<b>Selenium µg</b>	0.67	6.0
<b>Sodium mg</b>	13.4	40.2
<b>Zinc mg</b>	0.34	1.0
<b>OTHER</b>		
<b>Choline mg</b>	4.7	33.5
<b>Taurine mg</b>	N/S	8.0
<b>Nucleotides mg</b>	N/S	3.35
<b>Inositol mg</b>	2.68	26.8
<b>L-carnitine mg</b>	0.8***	N/S

See the notes on next page.

Note: Where regulations do not state a minimum amount, they are not considered a requirement. In these cases, the average stated for currently available first infant milks does not include those milks where the ingredient is not present.

N/S = not stated

- \* Vitamin E: 0.5mg/g of polyunsaturated fatty acids expressed as linoleic acid as corrected for the double bonds but in no case less than 0.5mg/100kcal, maximum 5.0mg/100kcal.
- \*\* For products manufactured from soya protein isolates or in a mixture with cows' milk, minimum and maximum values for iron for infant formula are 0.45mg and 2.0mg respectively, and for follow-on formula 0.9mg and 2.5mg respectively. For phosphorus, minimum and maximum values for both infant and follow-on formula are 30mg and 100mg respectively.
- \*\*\* The L-carnitine concentration is only specified for formula containing protein hydrolysates or soya protein isolates.

## Expert opinion on infant formula composition

In 2014 the European Food Safety Authority (EFSA) reviewed the essential composition of infant formula and follow-on formula (EFSA, 2014). EFSA independently analysed all the evidence available to determine the necessary, and unnecessary, components of infant formula, and considered other key issues related to ensuring products are safe and appropriate. Table 2 summarises some of the key points that EFSA made about infant formula (EFSA, 2014).

**TABLE 2**  
**Key points made by EFSA about infant formula**

Nutrient/component	EFSA (2014) opinion/comment
<b>Can formula milk be 'close' to breastmilk?</b>	<p>Breastmilk is the preferred food for all healthy infants. Whereas the composition of infant formula remains stable over time, breastmilk composition changes continuously and therefore infant formula cannot imitate breastmilk.</p> <p>Human milk composition can provide guidance on the composition of formula, but compositional similarity to human milk is not the only determinant or indicator of safety and nutritional suitability of formula.</p> <p>The mere presence of a substance in human milk does not necessarily indicate a specific benefit of this substance for the infant, nor do the concentrations of nutrients in human milk necessarily reflect infants' dietary requirements because they may mirror maternal intakes rather than infants' needs, or because absorption efficiency of certain nutrients differ between breastmilk and formula.</p> <p>Infant formula cannot imitate breastmilk with respect to its energy and protein content.</p>

Nutrient/component	EFSA (2014) opinion/comment
<p><b>Can formula milk be 'close' to breastmilk?</b></p>	<p>The structure of about 200 human milk oligosaccharides has been identified. The production of oligosaccharides is genetically determined and the individual pattern of oligosaccharides differs between women. The oligosaccharides of human milk are considered to be one of the principal growth factors, for example, for <i>Bifidobacteria</i> in the infant gut and are responsible for the composition of the gut microbiota found in breast-fed infants.</p> <p>There is no evidence for health benefits from the addition of prebiotic oligosaccharides (GOS/FOS) to infant or follow-on formula.</p>
<p><b>Is it safe to add ingredients that are not needed, or in amounts higher than necessary?</b></p>	<p>Nutrients and substances should be added to formulae for infants <i>only</i> in amounts that serve a nutritional or other benefit. The addition in amounts higher than those serving a nutritional or other benefit or the inclusion of unnecessary substances in formulae <b>may put a burden on the infant's metabolism or on other physiological functions</b>, as substances which are not used or stored have to be excreted.</p>
<p><b>Are there risks when nutrients are added at maximum amounts?</b></p>	<p>There is a lack of studies designed to investigate the short- or long-term health consequences of consumption of formulae containing the currently permitted maximum amounts of nutrients in infant formula.</p>
<p><b>Ingredients added to infant formula which are <u>unnecessary</u></b></p>	<ul style="list-style-type: none"> <li>• Arachidonic acid (ARA)</li> <li>• Eicosapentaenoic acid (EPA)</li> <li>• Non-digestible oligosaccharides (prebiotics. GOS/FOS mixtures)</li> <li>• Probiotics</li> <li>• Synbiotics (a mix of prebiotics and probiotics)</li> <li>• Chromium</li> <li>• Fluoride</li> <li>• Taurine</li> <li>• Nucleotides</li> <li>• Phospholipids as a source of long-chain polyunsaturated fatty acids instead of triacylglycerols</li> <li>• Triacylglycerols with palmitic acid predominantly esterified in the <i>sn</i>-2 position</li> </ul>

Source: EFSA, 2014.

## Which products are discussed in this resource?

**TABLE 3**  
**Infant milk products discussed in this resource**

Category of infant formula	Names of infant formula included in this category
<b>Infant formula suitable from birth (cows' milk based)</b>	Aptamil 1 First Milk Aptamil Profutura 1 First Infant Milk Bebivita First Infant Milk Castlemil First Infant Milk Cow & Gate 1 First Infant Milk Hipp Organic Combiotic First Infant Milk Holle Organic Infant Formula 1 Kendamil 1 First Infant Milk Kendamil Mehadrin First Infant Milk Mamia First Infant Milk Similac First Infant Milk SMA Pro First Infant Milk SMA Wysoy
<b>Infant formula suitable from birth (goats' milk based)</b>	Holle Organic Infant Goat Milk Formula 1 Kabrita Gold 1 NANNYcare First Infant Milk
<b>Infant formula marketed for hungrier babies, suitable from birth (cows' milk based)</b>	Aptamil Hungry Milk Cow & Gate Infant Milk for Hungrier Babies Hipp Organic Combiotic Hungry Infant Milk SMA Extra Hungry
<i>These milks are not recommended.</i>	

For each milk we summarise the key features of the milk, what claims have been made for it recently by the manufacturer, and its cost relative to other products. A summary of the costs of milks highlighted in this report can be found in the Appendix. A full summary of the costs of all infant milks on the UK market as of July 2017 can be found on the website [www.firststepsnutrition.org](http://www.firststepsnutrition.org).

## References

EFSA (2014). *Scientific opinion on the essential composition of infant and follow-on formulae*. Parma, Italy: European Food Safety Authority. Available at <http://www.efsa.europa.eu/en/efsajournal/pub/3760>

# Aptamil 1 First Milk

Aptamil 1 First Milk is sold in 900g tins of powder and as 70ml, 200ml and 1 litre RTF (ready-to-feed) bottles. The claims we discuss for this product were made on the previous Aptamil Professional website which has now been superseded by a new website covering all Aptamil and Cow & Gate brand products [www.eln.nutricia.co.uk](http://www.eln.nutricia.co.uk).

## Key components of this product

**TABLE 4**  
**Key components of Aptamil 1 First Milk**

Aptamil 1 First Milk		
	<b>Powdered infant milk</b>	<b>RTF (ready-to-feed) 1 litre bottle</b>
<b>Composition</b>	Meets current regulations for infant formula composition	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	Prebiotics, nucleotides, taurine, arachidonic acid	Prebiotics, nucleotides, taurine, arachidonic acid
<b>Allergens</b>	Cows' milk, soya, fish	Cows' milk, soya, fish
<b>Protein source</b>	Cows' milk	Cows' milk
<b>Whey:casein ratio</b>	60:40	60:40
<b>Fats used</b>	Palm, rapeseed, coconut and sunflower oils Fish and algal oils (single cell oils)	Palm, rapeseed, coconut and sunflower oils Fish and algal oils (single cell oils)
<b>Suitability</b>	Not suitable for vegetarians Halal approved	Not suitable for vegetarians
<b>Cost per 100ml</b>	17p	37p
<i>For comparison:</i> <b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on powdered formulation)	10p	10p

## Claims made about Aptamil 1 First Milk

### ***Contains Pronutra+, our unique blend of ingredients***

Pronutra+ is the name given by the manufacturer for a number of ingredients added to Aptamil brand formula milks that includes galacto- and fructo-oligosaccharides (GOS/FOS), long chain polyunsaturated fatty acids, and nucleotides.

**... our patented blend of Galacto- and Fructo-oligosaccharides (GOS/FOS) (9:1), which emulates the natural effect of breastmilk prebiotic oligosaccharides**

No specific references are used in support of this statement, but the website goes on to make further claims for the blend of GOS/FOS used in their infant formula milks:

**All Aptamil infant formulas contain a patented blend of GOS/FOS. Our blend of GOS/FOS (9:1) is shown to reduce the risk of allergies up to 5 years.**

and

**Babies fed a formula containing GOS/FOS have been shown to have a lower incidence of recurrent upper respiratory tract infections, diarrhoea, infections requiring antibiotics and atopic dermatitis than those fed a formula without GOS/FOS.**

References are given to studies sponsored by the manufacturer (Numico, now Danone) to support these claims. Those by Arslanoglu et al (2007) and Moro et al (2006) consider the impact of oligosaccharides in infant formula on infections and atopic dermatitis in the first 6 months of life, within the same cohort of infants at risk of atopy. Both studies reported reduced incidence of their outcome measures within the test groups compared to the control groups. The milk used in both the test and control groups was a formula containing extensively hydrolysed proteins supplemented with oligosaccharides, but Aptamil 1 First Milk does not contain hydrolysed proteins.

Another reference given to support the claims above is a study by Bruzzese et al (2009), also sponsored by Numico, which looked at the impact of infant formula milk supplemented with GOS/FOS at 0.4mg/100ml on the incidence of intestinal and respiratory infections in a cohort of healthy term infants. This study reported a lower incidence of gastroenteritis and fewer children receiving multiple courses of antibiotics within a year for the group receiving supplemented formula. The study was reviewed by EFSA (2014) as part of the evidence that led to their conclusion that there was insufficient evidence to support claims made about prebiotics in infant formula.

**These benefits have been shown to last for up to 5 years.**

The reference given to support this claim, Arslanoglu et al (2012), is a follow-up study from the previous Arslanoglu et al, 2007 study. The authors followed up some of the original cohort to 5 years, but only 42 children remained in the intervention group from the 102 original completers. The original study was reviewed and discounted as evidence by EFSA reviews of efficacy of benefit from oligosaccharide addition to infant formula.

**Bottlefed babies given a formula containing GOS/FOS also have softer stools than those fed a formula without GOS/FOS.**

Evidence is suggested to come from a study by Moro et al (2002), undertaken at the Numico (Danone) Research Centre. 90 healthy term infants were allocated to receive formula milk supplemented with oligosaccharides at a concentration of 0.4g/100ml or 0.8g/100ml or placebo, over a period of 28 days. The infant formula used in this study was different to

Aptamil 1 First Milk as it had higher protein and fat contents and a different range of added ingredients. The infants receiving the formula milk supplemented with oligosaccharides showed a dose-dependent increase in the amount of *Bifidobacteria* in stools. However, this does not mean that there is any clinical benefit to the addition of oligosaccharides.

It is disappointing that these claims are still being made for prebiotics when EFSA has repeatedly denied permission for a health claim based on the use of prebiotics in infant formula and follow on formula.

EFSA (2014) states that:

*“ ... there is no evidence for health benefits from the addition of prebiotic oligosaccharides (GOS/FOS) to infant or follow-on formula.”*

### **Increased levels of LCPs (long chain polyunsaturated fatty acids)**

Aptamil 1 First Milk currently contains 0.01g DHA/100ml (0.32% fatty acids). It is not clear from the website what ‘*increased levels*’ of LCPs relates to, as no indication is given as to whether previous formulations of this milk contained lower levels of DHA. This amount of DHA is within the range that all milks will have to contain by law when new regulations are brought in from July 2016. (Manufacturers are given a period of grace of several years from a change in regulation to update their products in line with any new recommendations.)

***LCPs at increased levels that are scientifically recognised by the European Food Safety Authority (EFSA) to support normal visual development and have a ratio of AA:DHA that meets consensus recommendations.***

The references given to support this statement are Koletzko et al (2008), and an EFSA opinion paper (EFSA, 2009).

The review article by Koletzko et al (2008) summarises the evidence for the role of DHA and AA in maternal and term infant nutrition as well as infant development, and makes recommendations concerning their inclusion in the infant diet. The review concluded that the fetus and neonate should receive LC-PUFA in amounts sufficient to support optimal visual and cognitive development. It recommended breastfeeding as the best way to achieve this and that if breastfeeding were not possible, formula supplemented with DHA and AA should be used. DHA supplementation should be at levels between 0.2 and 0.5 weight percent of total fat, with the minimum amount of AA equivalent to the contents of DHA.

In 2009, EFSA approved the claim that “*DHA contributes to the visual development of infants*” (EFSA, 2009). However, the validity of this claim is still debated, as it is argued that visual acuity develops slowly during the early years of life and early observations of visual measurements in infants are not predictive of later visual functions. Studies would have to follow children for seven years or longer to see if small clinical changes observed in early life had any real impact (Chambers et al, 2013).

The EFSA *Scientific opinion on the essential composition of infant and follow-on formulae* (EFSA, 2014) presents a useful summary of all the evidence relating to fatty acids in human and artificial milks, but concludes that, whilst they still believe that DHA should be added to

infant and follow-on formulae in similar amounts as are present in breastmilk as a 'prudent measure':

*"The panel notes there is no convincing evidence that the addition of LCPUFA to IF or FoF has any benefits beyond infancy on any functional outcomes."*

**... contains nucleotides, these form the building blocks of every cell in the body, including those of the immune system.**

Whilst no direct claims are made for the addition of nucleotides, Aptamil support their addition by reference to Pickering et al (1998) and Yau et al (2003).

Both of these clinical trials looked at the immune response of infants fed formula milk supplemented with nucleotides at 7.2mg/100ml. The trials had conflicting results. Yau et al (2003) reported that, at 8-28 weeks, infants fed the supplemented formula were shown to have a 25.4% lower risk of diarrhoea and higher concentrations of serum IgA throughout the study than infants fed the control formula. Both groups had a similar antibody response to hepatitis B immunisation and similar incidence of lower respiratory tract infections, whilst the risk of upper respiratory tract infections was 1.13 times higher in the group fed supplemented formula. Pickering et al (1998) reported that, compared to the control group, at 7 months of age the supplemented group had significantly higher H influenzae type b and diphtheria humoral antibody responses. The antibody responses to tetanus and polio virus were not enhanced by nucleotide fortification.

The fact that the formulas used in the trials were different to Aptamil 1 First Milk (which contains 3.2mg nucleotides/100ml), as well as the differences between outcomes in trials makes it difficult to see how this evidence might support the use of nucleotides in this infant formula. Both trials used milks supplemented with nucleotides at 7.2mg/100ml. Aptamil 1 First Milk is supplemented with nucleotides at 3.2mg/100ml. These studies were reviewed by EFSA (2014) and not accepted as evidence that nucleotides are beneficial.

EFSA (2014) state that:

*"there is no necessity to add nucleotides to infant or follow-on formula."*

### **Our conclusion**

Aptamil 1 First Milk meets the required compositional criteria for infant formula.

This formula contains a similar range of unnecessary added ingredients to some other brands on the market. However, in our opinion, there is no convincing evidence presented for the claims being made for these unnecessary ingredients, and claims are not supported by recent evidence reviews by expert bodies.

The differences between this formula and Aptamil Profutura 1 First Infant Milk do not warrant the price differential between the products.

### **References**

- Arslanoglu S, Moro GE, Boehm G (2007). Early supplementation of prebiotic oligosaccharides protects formula-fed infants against infections during the first 6 months of life. *Journal of Nutrition*; 137: 2420-24.
- Arslanoglu S, Moro GE, Boehm G et al (2012). Early neutral prebiotic oligosaccharide supplementation reduces the incidence of some allergic manifestations in the first 5 years of life. *Journal of Biological Regulators and Homeostatic Agents*; 26, 3: 49-59.
- Bruzzese E, Volpicelli M, Squeglia V et al (2009). A formula containing galacto- and fructo-oligosaccharides prevents intestinal and extra-intestinal infections: an observational study. *Clinical Nutrition*; 28, 2: 156-61.
- Chambers WA, Murphy D, Rodriguez W, Sun H (2013). Why any analysis of infant formula effects on infant visual acuity performed before the age of 7 years is not likely predictive. *Pediatrics*; 131, 1 e262-e272; 11 February 2013.
- EFSA (2014). *Scientific opinion on the composition of infant and follow-on formulae*. Parma, Italy: European Food Safety Authority. Available at <http://www.efsa.europa.eu/en/efsajournal/pub/3760>
- European Food Safety Authority Panel on Dietetic Products, Nutrition and Allergies (2009). Scientific opinion: DHA and ARA and visual development. *EFSA Journal*; 941: 1-14.
- Koletzko B, Lien E, Agostoni C et al (2008). The roles of long-chain polyunsaturated fatty acids in pregnancy, lactation and infancy: review of current knowledge and consensus recommendations. *Journal of Perinatal Medicine*; 36, 1: 5-14.
- Moro G, Arslanoglu S, Stahl B et al (2006). A mixture of prebiotic oligosaccharides reduces the incidence of atopic dermatitis during the first six months of age. *Archives of Disease in Childhood*; 91: 814-19.
- Moro G, Minoli I, Mosca M et al (2002) Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *Journal of Pediatric Gastroenterology and Nutrition*; 34, 3: 291-95.
- Pickering LK, Granoff DM, Erickson JR et al (1998). Modulation of the immune system by human milk and infant formula containing nucleotides. *Pediatrics*; 101 (2): 242-49.
- Yau K, Huang C, Chen W et al (2003). Effect of nucleotides on diarrhea and immune responses in healthy term infants in Taiwan. *Journal of Pediatric Gastroenterology and Nutrition*; 36: 37-43.

# Aptamil Profutura 1 First Infant Milk

Aptamil Profutura is available in 800g packs of powder, and 70ml and 200ml RTF bottles. Table 4 gives the data for the powder and 200ml RTF formula. The claims we discuss for this product were made on the previous Aptamil Professional website which has now been superseded by a new website covering all Aptamil and Cow & Gate brand products [www.eln.nutricia.co.uk](http://www.eln.nutricia.co.uk).

## Key components of this product

**TABLE 5**  
**Key components of Aptamil Profutura 1 First Infant Milk**

Aptamil Profutura 1 First Infant Milk		
	Powdered infant milk	RTF (ready-to-feed) 200ml bottle
<b>Composition</b>	Meets current regulations for infant formula composition	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	Prebiotics, nucleotides, taurine, phospholipids, arachidonic acid	Prebiotics, nucleotides, taurine, phospholipids, arachidonic acid
<b>Allergens</b>	Cows' milk, soya, egg, fish	Cows' milk, soya, egg, fish
<b>Protein source</b>	Cows' milk	Cows' milk
<b>Whey:casein ratio</b>	60:40	60:40
<b>Fats used</b>	Egg phospholipid Anhydrous milk fat, rapeseed oil, sunflower oil, coconut oil  Fish oil and algal oil (single cell oils)	Egg phospholipid Anhydrous milk fat, rapeseed, corn, coconut and sunflower oil  Fish oil and algal oil (single cell oils)
<b>Suitability</b>	Not suitable for vegetarians Halal approved	Not suitable for vegetarians
<b>Cost per 100ml</b>	21p	50p
<i>For comparison:</i> <b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on a powdered formulation)	10p	10p

## What claims are made for this infant formula?

### ***Egg lipid has been added.***

This is the first infant formula for healthy infants in the UK that contains egg lipid and which must therefore include egg as an allergen on the label (to add to the other allergens present: cows' milk, fish, milk and soya).

Danone, the manufacturer of this milk, have said in a letter to First Steps Nutrition Trust that:

*“The egg phospholipid in Aptamil Profutura milk RTF is sourced from egg yolk. We are confident that the egg phospholipid does not pose a risk of causing an allergic reaction and our supplier guarantees that the level of egg protein in the egg phospholipid is <0.001ppm per mg.”*

Apparently of the 11mg/100ml docosahexaenoic acid (DHA) present in the milk, 2.2mg comes from egg phospholipid. The remainder is provided by the fish oil and algal oils used.

### **Is there any benefit to adding phospholipids to infant formula?**

The Aptamil Professional website claims that Aptamil Profutura First Infant Milk:

*“Contains phospholipid-bound LCPs inspired by the complex structure of fats within breastmilk”.*

The EFSA 2014 opinion on the essential composition of infant formula stated that, whilst there are phospholipids naturally present in breastmilk:

*“There is no convincing evidence for a beneficial effect of using LCPUFA supplied as phospholipids in infant formula.”*

The manufacturer currently makes no claims specifically about the egg lipid as a source of LCPUFA in this formula, simply stating it is present, so it is not clear why they are using this as a partial source of DHA.

### ***Increased levels of LCPs (long chain polyunsaturated fatty acids).***

The manufacturers claim that Aptamil Profutura 1 First Infant Milk has increased levels of LCPs. Aptamil 1 First Milk currently contains 0.01g DHA/100ml, and Aptamil Profutura 1 First Infant Milk contains 0.011g/100ml (1/1000<sup>th</sup> g more per 100ml). This is a minimal change that is likely to be insignificant once compositional differences and compositional degradations within infant milk products are considered. These amounts are within the range that all milks will have to contain by law when new regulations come into force from July 2016. (Manufacturers are given a period of grace of three years from a change in regulation to update their products in line with any new recommendations.)

The manufacturers claim on the website that Aptamil Profutura 1 First Infant Milk:

*“Contains our highest levels of DHA to support normal visual development”.*

They support this claim with papers by Koletzko et al (2008), Willatts et al (1998), and an EFSA paper from 2009. All this data was reviewed by EFSA (2014) and they concluded that: *“The panel notes there is no convincing evidence that the addition of LCPUFA to IF or FoF has any benefits beyond infancy on any functional outcomes.”*

Furthermore, the claims do not make it clear that an increased amount of LCPs will not be of any greater benefit than the amount already used.

### ***The product uses anhydrous milk fat.***

The Aptamil Professional website claims that the product:

*“Contains natural milk fat to provide a better profile of fatty acids.”*

The website also claims:

*“In breastmilk, when palmitic acid is in the beta position it has certain benefits for babies, including better fat absorption, easy digestion, softer stools and reduced constipation.”*

Anhydrous milk fat is simply milk fat (butter fat) that has had all water removed. The references given to support this type of fat having properties which may aid fat and calcium absorption come from Jensen et al (1978), Carnielli et al (1996) and Kennedy et al (1999). The Jensen et al (1978) study reports on the variable composition of human milk in relation to maternal diet, and highlights the lack of reliable data at that time on milk composition. This paper offers no support for the statements made.

The studies by Carnielli et al (1996) and Kennedy et al (1999) have previously been used to support claims that use of *synthetic triglycerides* with a higher proportion of palmitate in the *sn-2* position improves fat and calcium absorption, but they are now being used to support the same claims for the use of *anhydrous milk fat*, which has a greater proportion of palmitate in the *sn-2* position than the vegetable oils commonly used in infant formula milks. Both of these studies examined the impact on fat and calcium absorption when infants were fed formula containing higher proportions of palmitic acid in the *sn-2* position than are present in Aptamil Profutura 1 First Infant Milk as a result of the use of anhydrous milk fat. The Kennedy et al study also reported that a number of parents reported concern about runny stools after feeding formula containing high *sn-2* palmitate.

The recent EFSA *Scientific opinion on the essential composition of infant and follow-on formulae* (2014) concluded that there was no convincing evidence for a beneficial effect of the use of palmitic acid predominantly esterified in the *sn-2* position in infant or follow-on formula.

### **Other claims made about Aptamil Profutura 1 First Infant Milk**

The following claims are made about Aptamil Profutura 1 First Infant Milk on the website or in print media.

### ***“Nutritionally closer to breastmilk than any other brand”***

It is important to note that EFSA (2014) clearly state in their opinion that:

*“Infant formula cannot imitate breast-milk with respect to its energy and protein content.”*

Aptamil Profutura support this claim with a chart comparing their milk with others, and base the claim on the added phospholipid and the fat composition which we have already reviewed above. They claim that:

***LCPs in breastmilk come in two different forms: phospholipid-bound and triglyceride-bound.***

The reference given to support this, Harzer et al (1983), looks at changing patterns of human milk lipids in the course of lactation, and clearly makes the point that milk composition changes as lactation progresses, and in mothers in different areas. It concludes that, because there are strong intra-individual differences in the composition of breastmilk, care should be taken when taking samples of breastmilk for analysis. This study does not in any way support a feed of consistent composition, or this milk, as being ‘close to breastmilk’. Just because manufacturers manipulate fat composition to mimic breastmilk components does not make it ‘closer to breastmilk’ in function.

The statement that this new milk is ‘closer to breastmilk’ is therefore not supported by any evidence provided in these references, nor in any of those references given to support fatty acid profiles of milks reviewed earlier.

The UK Guidance Notes (Department of Health, 2013) which explain the current regulations on infant formula, state that:

*“Non-mandatory text or pictures on infant formula and follow-on formula labelling must not make reference to ‘breastmilk’, ‘breastfeeding’, ‘moving on from breastfeeding’ or ‘closer to/inspired by breastmilk’. Use of such terms would not comply with Regulation 17 (2) or 18(2).”*

In our opinion, infant formula companies should not make claims that any infant formula is ‘closer to breastmilk’ on websites or in marketing materials, as this is not allowed on labels.

The website also makes the following claim about Aptamil Profutura First Infant Milk:

***Our innovative formulation also contains nucleotides, these form the building blocks of every cell in the body, including those of the immune system.***

Two references are given to support this: Pickering et al (1998) and Yau et al (2003). Both of these clinical trials looked at the immune response of infants fed formula milk supplemented with nucleotides at 7.2mg/100ml. Aptamil Profutura 1 First Infant Milk contains 3.2mg nucleotides/100ml so these studies do not directly relate to this formula. The trials had conflicting results.

Yau et al (2003) reported that, at 8-28 weeks, infants fed the supplemented formula were shown to have a 25.4% lower risk of diarrhoea and higher concentrations of serum IgA throughout the study than infants fed the control formula. Both groups had a similar antibody response to hepatitis B immunisation and similar incidence of lower respiratory tract infections, whilst the risk of upper respiratory tract infections was 1.13 times higher in the group fed supplemented formula. Pickering et al (1998) reported that, compared to the

control group, at 7 months of age, the supplemented group had significantly higher H influenzae type b and diphtheria humoral antibody responses. The antibody responses to tetanus and polio virus were not enhanced by nucleotide fortification. The fact that the formulas used in the trials were different to Aptamil Profutura 1 First Infant Milk, and the differences between outcomes in trials makes it difficult to interpret how this evidence might support the use of nucleotides in infant formula.

EFSA (2014) has stated that:

*“Taking into account the lack of convincing evidence for a benefit of the addition of nucleotides to IF and/or FOF, the Panel considers that there is no necessity to add nucleotides to IF or FOF.”*

***Contains our patented blend of GOS/FOS (9:1) bringing the intestinal microflora closer to that of a breastfed infant***

The manufacturers provide one reference from Moro et al (2002) to support this statement. In this study, 90 healthy term infants were randomly allocated to receive formula milk supplemented with oligosaccharides at a concentration of 0.4g/100ml, or 0.8g/100ml, or placebo, over a period of 28 days. The formulas used in this study have a different composition to Aptamil Profutura 1 First Infant Milk as they were higher in protein, lower in fat, had a different range of added ingredients and lower osmolality. The infants receiving the formula milk supplemented with oligosaccharides in this study showed a dose-dependent increase in the amount of *Bifidobacteria* in stools, but this alone does not provide evidence of clinical impact.

It is disappointing that this claim is still being made for prebiotics when EFSA has repeatedly denied permission to make a claim based on the use of prebiotics in infant formula and follow-on formula.

EFSA (2014) states that:

*“... there is no evidence for health benefits from the addition of prebiotic oligosaccharides (GOS/FOS) to infant or follow-on formula.”*

### **Our conclusion**

Aptamil Profutura 1 First Infant Milk meets the required compositional criteria for infant formula.

In our opinion, this formula offers no advantage over any other first infant milk. The very high cost of this milk, the increased range of potential allergens included, and the high number of unnecessary ingredients could be seen as making this a less good choice.

EFSA (2014) clearly state: *“Nutrients and other substances should be added to formulae for infants only in amounts that serve a nutritional or other benefit. The addition in amounts higher than those serving a benefit, or the inclusion of unnecessary substances in formulae puts a burden on the infant’s metabolism and/or physiological functions as substances which are not used or stored have to be excreted.”*

Using 70ml RTF bottles of Aptamil Profutura 1 First Infant Milk to feed an infant in the first week of life would cost over £100. (For a list of the cost of various types of infant milks, see Appendix 1)

The differences between this formula and Aptamil 1 First Milk do not warrant the price differential between the products.

## References

Carnielli VP, Luijendijk IH, Van Godoever J et al (1996). Structural position and amount of palmitic acid in infant formulas: effects on fat, fatty acid, and mineral balance. *Journal of Pediatric Gastroenterology and Nutrition*; 23: 533-60.

Department of Health (2013). *DH Guidance Notes on the Infant Formula and Follow-on Formula Regulations 2007 (as amended)*. Available at: [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/204314/Infant\\_formula\\_guidance\\_2013\\_-\\_final\\_6\\_March.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/204314/Infant_formula_guidance_2013_-_final_6_March.pdf)

EFSA (2014). *Scientific opinion on the essential composition of infant and follow-on formulae*. Parma, Italy: European Food Safety Authority. Available at <http://www.efsa.europa.eu/en/efsajournal/pub/3760>

European Food Safety Authority Panel on Dietetic Products, Nutrition and Allergies (2009). Scientific opinion: DHA and ARA and visual development. *EFSA Journal*; 941: 1-14.

Harzer G, Haug M, Dieterich I, Gentner PR (1983). Changing patterns of human milk lipids in the course of the lactation and during the day. *American Journal of Clinical Nutrition*; 37 (4): 612-21.

Jensen R, Hagerty M, McMahon K (1978). Lipids of human milk and infant formulas: A review. *American Journal of Clinical Nutrition*; 31: 990-1016.

Kennedy K, Fewtrell MS, Morley R et al (1999). Double-blind, randomized trial of a synthetic triacylglycerol in formula-fed term infants: effects on stool biochemistry, stool characteristics, and bone mineralization. *American Journal of Clinical Nutrition*; 70: 920-27.

Koletzko B, Lien E, Agostoni C et al (2008). The roles of long-chain polyunsaturated fatty acids in pregnancy, lactation and infancy: review of current knowledge and consensus recommendations. *Journal of Perinatal Medicine*; 36, 1: 5-14.

Moro G, Minoli I, Mosca M et al (2002). Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *Journal of Pediatric Gastroenterology and Nutrition*; 34, 3: 291-95.

Pickering LK, Granoff D, Erickson J et al (1998). Modulation of the immune system by human milk and infant formula containing nucleotides. *Pediatrics*; 101 (2): 242-49.

Willatts P, Forsyth JS, DiModugno MK et al (1998). Effect of long-chain polyunsaturated fatty acids in infant formula on problem solving at 10 months of age. *Lancet*; 352: 688-91.

Yau K, Huang C, Chen W et al (2003). Effect of nucleotides on diarrhea and immune responses in healthy term infants in Taiwan. *Journal of Pediatric Gastroenterology and Nutrition*; 36: 37-43.

# Bebivita First Infant Milk

---

Bebivita is sold at LIDL stores only in the UK and is marketed in 500g packets only.

## Key components of this product

**TABLE 6**  
**Key components of Bebivita First Infant Milk**

Bebivita First Infant Milk	
	<b>Powdered infant milk</b>
<b>Composition</b>	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	
<b>Allergens</b>	Cows' milk, soya
<b>Protein source</b>	Cows' milk
<b>Whey:casein ratio</b>	60:40
<b>Fats used</b>	Palm, rapeseed, sunflower oils
<b>Suitability</b>	Not suitable for vegetarians Not Halal approved
<b>Cost per 100ml<sup>1</sup></b>	10p
<i>For comparison:</i>	10p
<b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on powdered formulation)	

This milk has been marketed in UK LIDL stores since September 2016 There is currently no marketing information in the UK for this product but we will update this section if this becomes available.

### Our conclusion

Bebivita First Infant Milk meets the required compositional criteria for infant formula. This milk contains no unnecessary added ingredients, but is not suitable for families seeking a vegetarian or halal product. This milk is currently one of the least expensive powdered infant formulas on the market.

# Castlemil Infant Milks

These cows' milk based infant formula milks are manufactured by Castlemil and are currently only available on-line from a limited number of retailers. The products are available as powder formulation in 800g tins.

## Key components of this product

**Table 7 Key components of Castlemil Infant Milk**

Castlemil First Infant Milk	
	<b>Powdered infant milk</b>
<b>Composition</b>	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	Arachidonic acid, prebiotics, taurine, nucleotides
<b>Allergens</b>	Cows' milk, soya, fish
<b>Protein source</b>	Cows' milk
<b>Whey:casein ratio</b>	57:33
<b>Fats used</b>	Sunflower, rapeseed, palm, coconut, fish and algal oils (single cell oils) milk fat
<b>Suitability</b>	Not suitable for vegetarians Not Halal approved
<b>Cost per 100ml<sup>1</sup></b>	15p
<i>For comparison:</i> <b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on powdered formulation)	10p

## Claims made

No specific claims are made for these milks on the company website, however it does state that the milks are made to exacting standards using British Red Tractor farm assured whole milk from local dairies and produced by free range, grass fed cows. The whole milk used in the products, after drying, accounts for about 11%, 17% and 30% of the dry weight of the milk powders for infant, follow-on and toddler milks respectively.

# Cow & Gate 1 First Infant Milk

Cow & Gate 1 First Infant Milk is available in powdered form (900g), 70ml , 200ml and 1 litre RTF (ready-to-feed) bottles. The claims we discuss for this product were made on the previous Cow & Gate website for health professionals 'In Practice' which has now been superseded by a new website covering all Aptamil and Cow & Gate brand products [www.eln.nutricia.co.uk](http://www.eln.nutricia.co.uk).

## Key components of this product

**TABLE 8**  
**Key components of Cow & Gate 1 First Infant Milk**

Cow & Gate 1 First Infant Milk		
	Powdered infant milk	RTF (ready-to-feed) 200ml carton
<b>Composition</b>	Meets current regulations for infant formula composition	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	Prebiotics, nucleotides, taurine, arachidonic acid	Prebiotics, nucleotides, taurine, arachidonic acid
<b>Allergens</b>	Cows' milk, soya, fish	Cows' milk, soya, fish
<b>Protein source</b>	Cows' milk	Cows' milk
<b>Whey:casein ratio</b>	60:40	60:40
<b>Fats used</b>	Palm, rapeseed, coconut and sunflower oils Fish and algal oils (single cell oils)	Palm, rapeseed, coconut and sunflower oils Fish and algal oils (single cell oils)
<b>Suitability</b>	Not suitable for vegetarians Halal approved	Not suitable for vegetarians
<b>Cost per 100ml</b>	14p	38p
<i>For comparison:</i> <b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on a powdered formulation)	10p	10p

## Claims made about Cow & Gate 1 First Infant Milk

Cow & Gate do not make specific claims for their First Infant Milk on the product pages of their 'In Practice' website. They simply state some of the ingredients used and make a general claim that that these help ensure healthy growth and development:

***Cow & Gate first infant milk from newborn contains prebiotic oligosaccharides (OS), nucleotides, a fat blend that includes LCPs, calcium : phosphorus in a ratio that aids calcium absorption, and added vitamins, minerals and antioxidants, to help ensure healthy growth and development.***

No claims are made for any specific benefit associated with any of these components and the entire statement is supported by nine references.

### **Cow & Gate Infant Milk contains nucleotides.**

The two papers we think are being used to support the addition of nucleotides are Pickering et al (1998) and Maldonado Lozano et al (2001). Pickering et al studied two groups of healthy term infants that were fed either formula milk supplemented with nucleotides at 7.2mg/100ml or formula that was not supplemented. They reported no differences between the groups at 6 months of age, but at 7 months of age, the supplemented group had significantly higher H influenzae type b and diphtheria humoral antibody responses. The antibody responses to tetanus and polio virus were not enhanced by nucleotide fortification. Maldonado Lozano et al (2001) reviewed the literature on the impact of adding nucleotides to cows' milk based infant formula and concluded that their addition seems to produce a favourable effect on the immune function. Cow & Gate 1 First Infant formula contains 3.2mg nucleotides/100ml and is therefore not equivalent to the milk used in this study.

EFSA (2014) have concluded that:

*“Taking into account the lack of convincing evidence for a benefit of the addition of nucleotides to IF and/or FOF, the Panel considers that there is no necessity to add nucleotides to IF or FOF.”*

### **Cow & Gate Infant Milk contains LCPs.**

The three papers we believe are used to support the addition of LCPs are Willatts et al (1998), Birch et al (1998), and Birch et al (2000).

The Willatts et al (1998) study, funded by Milupa (Danone), considers long chain fatty acid supplementation in term infants through formula supplementation and impact on IQ. Cognitive behaviour was tested at 10 months by a three-step solution to uncovering and retrieving a hidden toy. The authors conclude that there may be some benefit of supplementation with long chain polyunsaturated fatty acids, but whether it is possible to measure cognitive behaviour at this age is debateable.

Birch et al (1998) reported that the supplementation of term infant formula with DHA or with DHA and AA during the first 4 months of life yields better sweep visual evoked potential acuity at 6, 17 and 57 weeks of age but not at 26 weeks when acuity development reaches a plateau.

Birch et al (2000) reported a statistically significant developmental age advantage for DHA and DHA+AA supplemented groups in the cognitive and motor subscales of the Bayley Scales of Infant Development compared to the unsupplemented group, but no such increase was observed for the language, psychomotor development index or behaviour rating scale. On the basis of their results, the authors suggest that early dietary supply of DHA was a dietary determinant of improved performance on the mental development index.

None of these conclusions are supported by EFSA (2014), who conclude that:

*“The panel notes there is no convincing evidence that the addition of LCPUFA to IF or FoF has any benefits beyond infancy on any functional outcomes.”*

### **Cow & Gate Infant Milk contains prebiotic oligosaccharides.**

The three papers we believe are referenced to support the addition of prebiotics were all sponsored by Numico (Danone).

Arslanoglu et al (2008) extends a previous study (also sponsored by Danone) in which healthy term infants at risk of atopy were fed either a hypoallergenic (hydrolysed) formula supplemented with GOS/FOS or the same formula with placebo of maltodextrins. The aim of this study was to evaluate whether or not protective effects against allergic manifestations and infections observed in an initial 6-month trial and attributed by the authors to GOS/FOS, extended as far as 2 years of life. The authors reported that at 2 years the cumulative incidences of atopic dermatitis, recurrent wheezing and allergic urticaria (itching) were lower in the group that received GOS/FOS compared to the placebo group, but it is difficult to attribute this solely to early formula use.

Bruzzese et al (2009) looked at the impact of standard infant formula supplemented with GOS/FOS on the incidence of intestinal and respiratory infections in a cohort of healthy, term infants. This study reported a lower incidence of gastroenteritis, and fewer children receiving multiple courses of antibiotics within a year for the group receiving supplemented formula.

Knol et al (2005) looked at whether or not standard infant formula supplemented with GOS/FOS was able to establish a bifido-dominant microflora, not only in numbers, but also with respect to the metabolic activity in the colon. After 6 weeks, the mean proportion of *Bifidobacteria* was significantly higher in the group given formula with GOS/FOS and the authors concluded that the addition of the prebiotic GOS/FOS mixture to an infant formula had a stimulating effect on the growth of *Bifidobacteria* and on the metabolic activity of the total intestinal flora.

All of these studies were included in the review by EFSA (2014) but were not accepted as evidence of benefit from the addition of GOS/FOS to infant formula.

Whilst no direct claims are made on the product pages of the Cow & Gate 'In Practice' website, the professional resources and bottle feeding pages place great emphasis on prebiotics and the following claims are made:

***... mums can encourage the development of beneficial bacteria in the gut by choosing a formula that contains ... Prebiotic oligosaccharide supplements: non-digestible carbohydrates that selectively stimulate the growth and activity of beneficial bacteria. Different formulas may contain different types and amounts of prebiotics.***

This statement is supported by reference to a clinical trial conducted by the Numico (Danone) Research in which 90 healthy term infants were randomly allocated to receive formula milk supplemented with oligosaccharides over a period of 28 days Group Moro et al (2002). The authors reported that after 28 days there was a dose-dependent increase in the number of *Bifidobacteria* in the stools of both groups receiving GOS/FOS, but this alone does not provide evidence of clinical benefit.

There is a lot of emphasis placed on the specific proportions and type of oligosaccharides used in Cow & Gate 1 First Infant Formula and the following claims are made:

***Possible benefits of the GOS/FOS blend include a reduction in:***

***Intestinal infections (including diarrhoea)***

***Recurrent respiratory tract infections***

***Infections requiring antibiotics***

***Allergic symptoms, including atopic dermatitis***

***This specific blend has been shown to have a number of benefits in infants:***

***Discourages the growth of harmful pathogens***

***Promotes healthy digestion, as indicated by softer more frequent stools***

***Improves feeding tolerance, with significantly less regurgitation and crying in connection with feeding***

***... reduced the incidence of allergic manifestations, such as atopic dermatitis, for up to 5 years in children with a predisposition for allergies.***

The particular blend of oligosaccharides used in Cow & Gate Infant Milks was subject to a wide range of clinical trials carried out by, or sponsored by, the then parent company Numico (now Danone).

The study by Knol et al (2005) which is outlined above is used again to support the claim that the specific blend of prebiotics used discourages the growth of harmful pathogens. However, this study looked at whether or not standard infant formula supplemented with GOS/FOS is able to establish a bifido-dominant microflora. It does not therefore directly support the claim concerning harmful pathogens.

Studies by Moro et al (2002) and (2006) are used to support claims related to increased stool softness and improved feeding tolerance, despite this being generally a subjective measure. Moro et al (2002) reported that stool softness was only increased in the group receiving formula supplemented at 0.8g/dL and not in those receiving 0.4g/100ml GOS/FOS. Cow & Gate 1 First Infant Milk is currently supplemented at a rate of 0.6g/100ml, but this infant formula is different to the one used in the studies as it has a lower protein and fat content, a different range of added ingredients and a higher osmolality.

The main outcome measure of the study by Moro et al (2006) was incidence of atopic dermatitis between groups fed formula supplemented or not supplemented with GOS/FOS.

However, this study was used to support the claim: *“Improves feeding tolerance, with significantly less regurgitation and crying in connection with feeding.”*

It is interesting to note that the Moro (2002) study outlined above actually reported a very different result:

*“The different diets did not influence the incidence of crying, regurgitation, or vomiting.”*

The website cites three papers by Arslanoglu et al in support of the claims made concerning atopic dermatitis and infections. These are a series of studies which follow the same cohort of children at risk of atopy from the first 6 months of life to 5 years. The original study by Arslanoglu et al (2007) reports on the impact of supplementation of hypoallergenic, hydrolysed whey-based formula milk with either 0.8g/100ml GOS/FOS or 0.8g/100ml maltodextrin as placebo on atopic dermatitis and allergy symptoms in the first 6 months of life. The Arslanoglu et al (2008) and Arslanoglu et al (2012) studies extend the previous (flawed) study published in 2007 to 2 years and 5 years respectively. At 5 years only 42 children remained in the intervention group from the 102 original completers.

The original study (Arslanoglu et al, 2007) was reviewed and discounted as evidence by earlier reviews of efficacy of benefit from oligosaccharide addition to infant formula and was also reviewed by EFSA (2014).

The claims for the addition of GOS/FOS have been made for many years, despite a clear statement from EFSA (2014) that:

*“There is insufficient evidence for beneficial effects on infant health of the oligosaccharides that have been tested to date in RCTs when added to infant or follow-on formula.”*

### **Our conclusion**

Cow & Gate 1 First Infant Milk meets the required compositional criteria for infant formula.

This milk contains a similar range of unnecessary added ingredients as other brands on the market. It offers no evidence-based advantage over any other first infant milk. The GOS/FOS blend used by Cow & Gate has been emphasised on the website, creating the impression that this is superior to those used by other manufacturers, but the evidence used to support the inclusion of any type or blend of prebiotic has been clearly refuted by EFSA.

This milk represents better value for money than some other brands. (For a list of the cost of various types of infant milks, see the Appendix.)

### **References**

Arslanoglu S, Moro G, Boehm G (2007). Early supplementation of prebiotic oligosaccharides protects formula-fed infants against infections during the first 6 months of life. *Journal of Nutrition*; 137: 2420-24.

Arslanoglu S, Moro GE, Boehm G et al (2012). Early neutral prebiotic oligosaccharide supplementation reduces the incidence of some allergic manifestations in the first 5 years of life. *Journal of Biological Regulators and Homeostatic Agents*; 26, 3: 49-59.

Arslanoglu S, Moro GE, Schmitt J et al (2008). Early dietary intervention with a mixture of prebiotic oligosaccharides reduces the incidence of allergic manifestations and infections during the first 2 years of life. *Journal of Nutrition*; 138, 6: 1091-95.

Birch E, Garfield S, Hoffman D et al (2000). A randomized controlled trial of early dietary supply of long-chain polyunsaturated fatty acids and mental development in term infants. *Developmental Medicine and Child Neurology*; 42: 174-81.

Birch EE, Hoffman DR, Uauy R et al (1998). Visual acuity and the essentiality of docosahexaenoic acid and arachidonic acid in the diet of term infants. *Pediatric Research*; 44: 201-09.

Bruzzese E, Volpicelli M, Squeglia V et al (2009). A formula containing galacto- and fructo-oligosaccharides prevents intestinal and extra-intestinal infections: an observational study. *Clinical Nutrition*; 28, 2: 156-61.

Knol J, Scholtens P, Kafka C et al (2005). Colon microflora in infants fed formula with galacto- and fructo-oligosaccharides: more like breast-fed infants. *Journal of Pediatric Gastroenterology and Nutrition*, 40, 36-42.

Maldonado Lozano J, Navarro J, Narbona E, Gil A (2001). The influence of dietary nucleotides on humoral and cell immunity in the neonate and lactating infant. *Early Human Development*; 65 (Suppl): S69-74.

Moro GE, Minoli I, Mosca M et al (2002). Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *Journal of Pediatric Gastroenterology and Nutrition*, 34, 291-95.

Moro G, Arslanoglu S, Stahl B et al (2006). A mixture of prebiotic oligosaccharides reduces the incidence of atopic dermatitis during the first six months of age. *Archives of Disease in Childhood*; 91: 814-19.

Pickering LK, Granoff DM, Erickson JR et al (1998). Modulation of the immune system by human milk and infant formula containing nucleotides. *Pediatrics*; 101 (2): 242-49.

Willatts P, Forsyth JS, DiModugno M et al (1998). Effect of long-chain polyunsaturated fatty acids in infant formula on problem solving at 10 months of age. *Lancet*, 352, 688-91.

# HiPP Organic Combiotic First Infant Milk

This milk is available as a powdered formulation in 800g powder packs and as 200ml and 90ml RTF (ready-to-feed). Table 8 gives details for the powder and 200ml RTF.

## Key components of this product

**TABLE 9**  
**Key components of HiPP Organic Combiotic First Infant Milk**

HiPP Organic Combiotic First Infant Milk		
	Powder	RTF (ready-to-feed) 200ml carton
<b>Composition</b>	Meets current regulations for infant formula composition	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	Prebiotics, arachidonic acid	Prebiotics, arachidonic acid,
<b>Allergens</b>	Cows' milk, fish	Cows' milk, soya
<b>Protein source</b>	Cows' milk	Cows' milk
<b>Whey:casein ratio</b>	60:40	60:40
<b>Fats used</b>	Rapeseed, palm, coconut and sunflower oils Fish and algal oils (single cell oils)	Palm, rapeseed and sunflower oils Fungal and algal oils (single cell oils)
<b>Suitability</b>	Not suitable for vegetarians Not halal approved	Not suitable for vegetarians Not halal approved
<b>Cost per 100ml</b>	14p	38p
<i>For comparison:</i> <b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on a powdered formulation)	10p	10p

## Claims made for this product

Hipp Organic have said that the claim that this product has the lowest protein content of all infant formula on the market has now been withdrawn, but the data card for this milk on the health professional website still includes a graph that clearly shows their milk with a lower protein content than other standard infant formula. We have included the claims made in marketing materials from Hipp in the past year in the discussion below.

**... the UK's first reduced protein infant formula, now with alpha-lactalbumin, making the protein level and profile closer to that found in breastmilk**

The reference given to support this statement, Nommsen et al (1991), is from an American study which examined factors associated with concentrations of energy-yielding nutrients in human milk. This study measured the protein in samples of milk from lactating women at different time points over a period of 12 months, but the protein profile was not examined. This study also showed that there was significant variation in breastmilk composition related to a number of maternal factors, and it is important to remember that breastmilk changes composition during and between feeds and over time. The statement that this infant formula has “a protein level and profile closer to breastmilk” is therefore not supported by this reference.

It is important to note that EFSA (2014) clearly state in their opinion that:

*“Infant formula cannot imitate breast-milk with respect to its energy and protein content.”*

The UK Guidance Notes (Department of Health, 2013) which explain the current regulations on infant formula, state that:

*“Non-mandatory text or pictures on infant formula and follow-on formula labelling must not make reference to ‘breastmilk’, ‘breastfeeding’, ‘moving on from breastfeeding’ or ‘**closer to/inspired by breastmilk**’. Use of such terms would not comply with Regulation 17 (2) or 18(2).”*

In our opinion, infant formula companies should not make claims that any infant formula is ‘closer to breastmilk’ on websites or in marketing materials as this is not allowed on labels.

**the UK's first reduced protein infant formula**

HiPP was the first manufacturer to introduce infant formula with a protein level of less than 2g/100kcal to the UK market. However, the difference is not sufficiently significant to differentiate it from other brands, and in terms of protein content per 100ml of infant formula most first formula now have comparable amounts, with the four most popular brands all having about 1.3g/100ml. (For full compositional data see the report *Infant Milks in the UK* at [www.firststepsnutrition.org](http://www.firststepsnutrition.org))

HiPP makes claims for its product based on the results of the HiPP-funded BeMIM study (Fleddermann et al, 2014). The aim of this study was to prove the safety and suitability of a reduced-protein formula for healthy term babies, and was not designed to either prove, or disprove, the theory that early intakes of protein and energy influence obesity risk in later life. The study compared outcomes related to the intakes of two formula, one with a protein

content of 1.3g/100ml, and one with a protein content of 1.5g/100ml. There are no first infant formula on the market with a protein content of 1.5g/100ml.

The claims made by HiPP for their formula are based on findings from this study and include:

#### **Adequate growth**

This is to be expected as the low protein test formula milk meets current compositional recommendations.

#### **Good tolerance and acceptance**

This is also to be expected, as the composition was similar to other formula.

#### **Protein intakes closer to breastfed infants**

The fact that protein intakes may be similar in breastfed and formula-fed infants does not suggest that this is associated with any clinical advantage. The estimate used for protein intakes of breastfed infants are based on breastmilk composition of 1.2g/100ml (1.7g/100kcal), reported by Nommsen et al (1991). The current estimate of protein content in breastmilk in the UK is 1.3g/100ml (Finglas et al, 2015).

The evidence presented by HiPP to promote their milks on the basis of reduced protein does not suggest that this infant formula has any clinical advantage over other whey-based first infant formula available on the market.

#### **Added LCPs omega 3 & 6 – Docosahexaenoic acid (DHA) and Arachidonic acid (AA), important for the development of the brain, nervous system and eyesight**

The reference given to support this statement is Koletzko et al (2008). This review article summarised the evidence for the role of DHA and AA in maternal and term infant nutrition as well as infant development and made recommendations concerning their inclusion in the infant diet. The review concluded that the fetus and neonate should receive LC-PUFA in amounts sufficient to support optimal visual and cognitive development. It recommended breastfeeding as the best way to achieve this and if breastfeeding were not possible, formula supplemented with DHA and AA should be used.

The EFSA *Scientific opinion on the essential composition of infant and follow-on formulae* (EFSA, 2014) presents a useful summary of all the evidence relating to fatty acids in human and artificial milks, but concludes that, whilst they believe that DHA should be added to infant and follow-on formulae in similar amounts as are present in breastmilk as a “*prudent measure*”, there is:

*“currently no conclusive evidence for any effects beyond infancy of DHA supplementation in any of the health outcomes studied.”*

#### **Added prebiotic oligosaccharides (galacto-oligosaccharides – GOS), extracted from lactose**

No specific claims are made for the addition of galacto-oligosaccharides on the health professional website. However, Ben et al (2008) is referenced to support the addition of GOS. This pilot study conducted in China in a relatively small sample of infants (with no intention to treat analysis attempted) looked at levels of *Bifidobacteria* and *Lactobacilli* and stool characteristics in infants fed either formula milk with or without GOS, those having human milk and formula with GOS, or those having human milk only. After three months, the intestinal *Bifidobacteria* and *Lactobacilli*, were somewhat higher in the group fed formula with GOS than in the group given unsupplemented formula acetic acid and fecal pH was decreased in infants fed with the GOS-formula or human milk. The formula used in this study was also a different type of formula to the HiPP formula described here that uses this reference to support its claims. This study was reviewed by EFSA (2014) which concluded that there was no significant evidence that the addition of prebiotics are beneficial to infant health.

EFSA (2014) state that:

*“there is no evidence for health benefits from the addition of prebiotic oligosaccharides (GOS/FOS) to infant or follow-on formula.”*

**Organic – made from milk produced by cows kept to organic standards and fed an organic diet, free from GM ingredients and chemical pesticides**

The organic status of HiPP Organic Combiotic First Infant Milk distinguishes it from most other infant milks on the UK market. However, it is important to note that there are limits on the level of any individual pesticide residue that may be present in infant formula and follow-on formula and specific upper limits for toxic pesticides.

HiPP Organic reference a study by Butler et al (2008) to support the statements that organic milk is higher in vitamin E, beta-carotene, lutein and zeaxanthine, but fails to mention what the comparison is. The Butler et al study, which compares concentrations of fatty acids and antioxidants in milk from high-input, conventional farms, low-input organic farms and low-input non-organic farms, reported that milk from the low-input organic *and* low-input non-organic farms had higher concentrations of conjugated linoleic and  $\alpha$ -linolenic acid,  $\alpha$ -tocopherol and carotenoids compared with milk from the high-input system. Milk composition differed significantly between the two low-input systems during the second half of the grazing period only, with milk from non-organic cows being higher in antioxidants and conjugated linoleic acid, and that from organic cows being higher in  $\alpha$ -linolenic acid. This does not necessarily translate into higher levels in infant formula made with organic milk where concentrations of micronutrients are regulated; nor does it suggest any clinical advantage of higher levels of micronutrients.

### **Our conclusion**

HiPP Organic Combiotic First Infant Milk meets all the required compositional criteria for infant formula.

In our opinion it offers no clinical advantage over any other first infant milk, but it has fewer unnecessary ingredients added than some other types of milk.

If families want to use an organic infant formula milk, this product offers better value for money than available alternatives. (For a list of the cost of various types of infant milks, see the Appendix.)

## References

Ben XM, Li J, Feng ZT et al (2008). Low level of galacto-oligosaccharide in infant formula stimulates growth of intestinal *Bifidobacteria* and *Lactobacilli*. *World Journal of Gastroenterology*; 14: 6564-68.

Butler G, Nielsen JH, Slots T et al (2008). Fatty acid and fat-soluble antioxidant concentrations in milk from high- and low-input conventional and organic systems: seasonal variation. *Journal of the Science of Food and Agriculture*; 88: 1431-41.

Department of Health (2013). *DH Guidance Notes on the Infant Formula and Follow-on Formula Regulations 2007 (as amended)*. Available at: [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/204314/Infant\\_formula\\_guidance\\_2013\\_-\\_final\\_6\\_March.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/204314/Infant_formula_guidance_2013_-_final_6_March.pdf)

EFSA (2014). *Scientific opinion on the composition of infant and follow-on formulae*. Parma, Italy: European Food Safety Authority. Available at <http://www.efsa.europa.eu/en/efsajournal/pub/3760>

Finglas PM, Rose MA, Pinchen HM et al (2015) *McCance and Widdowson's The Composition of Foods*, Seventh summary edition. Cambridge: Royal Society of Chemistry.

Fleddermann M, Demmelmair H, Grote V et al (2014). Infant formula composition affects energetic efficiency for growth: The BeMIM study, a randomized controlled trial. *Clinical Nutrition*; 33: 588-95.

Koletzko B, von Kreis R, Closa R et al (2009). Lower protein in infant formula is associated with lower weight up to age 2y: a randomized clinical trial. *American Journal of Clinical Nutrition*; 89: 1836-45.

Koletzko B, Lien E, Agostoni C et al (2008). The roles of long-chain polyunsaturated fatty acids in pregnancy, lactation and infancy: review of current knowledge and consensus recommendations. *Journal of Perinatal Medicine*; 36; 1: 5-14.

Nommsen LA, Lovelady CA, Heinig MJ et al (1991). Determinants of energy, protein, lipid and lactose concentrations in human milk during the first 12 mo of lactation: the DARLING Study. *American Journal of Clinical Nutrition*; 53 (2): 457-65.

# Holle Organic Infant Formula 1

This organic cows' milk based infant formula is available in specialist health food shops in powdered form in the UK. It is imported from Switzerland.

## Key components of this product

**TABLE 10**  
**Key components of Holle Organic Infant Formula 1**

Holle Organic Infant Formula 1	
	Powdered infant milk (400g box)
<b>Composition</b>	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	None
<b>Allergens</b>	Cows' milk
<b>Protein source</b>	Cows' milk
<b>Whey:casein ratio</b>	60:40
<b>Fats used</b>	Palm, rapeseed and sunflower oils
<b>Suitability</b>	Not suitable for vegetarians Not halal approved
<b>Cost per 100ml</b>	28p
<i>For comparison:</i> <b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on a powdered formulation)	10p
<b>Preparation instructions</b>	The instructions recommend that the formula is made up with water at 50°C which is not in line with UK recommendations. <b>We recommend that the formula is made up following UK guidance.</b>

## Claims made

No specific claims are made for this milk except that the milk used to prepare this infant formula is 'biodynamic.' Biodynamic farming shares many principles with organic farming but also has a spiritual-ethical dimension. Information about biodynamic farming can be found at [www.biodynamics.com](http://www.biodynamics.com).

# Holle Organic Infant Goat Milk Formula 1

This organic goats' milk based infant formula is available in specialist health food shops in powdered form in the UK. It is imported from Switzerland.

## Key components of this product

TABLE 11

Key components of Holle Organic Infant Goat Milk Formula 1

Holle Organic Infant Goat Milk Formula 1	
Powdered infant milk (400g box)	
<b>Composition</b>	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	None
<b>Allergens</b>	Goats' milk
<b>Protein source</b>	Goats' milk
<b>Whey:casein ratio</b>	14:83
<b>Fats used</b>	Palm, rapeseed and sunflower oils
<b>Suitability</b>	Suitable for vegetarians Not halal approved
<b>Cost per 100ml</b>	35p
<i>For comparison:</i> <b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on a powdered formulation)	10p
<b>Preparation instructions</b>	The instructions recommend that the formula is made up with water at 50°C, which is not in line with UK recommendations. <b>We recommend that the formula is made up following UK guidance.</b>

## Claims made

No specific claims are made for this milk, except that the milk used to prepare this infant formula is organic.

# Kabrita Gold 1

This goats' milk based formula is available in 800g tins in specialist food shops.

## Key components of this product

**TABLE 12**  
**Key components of Kabrita Gold 1**

Kabrita Gold 1	
	Powdered infant milk
<b>Composition</b>	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	Prebiotics, taurine
<b>Allergens</b>	Goats' milk, fish, soya
<b>Protein source</b>	Goats' milk
<b>Whey:casein ratio</b>	64:36
<b>Fats used</b>	Soybean, sunflower and palm kernel oil; with high sn-2 palmitic acid oil
<b>Suitability</b>	Not suitable for vegetarians Not halal approved
<b>Cost per 100ml</b>	33p
<i>For comparison:</i> <b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on a powdered formulation)	11p

## Claims made

***Is naturally easy to digest (due to a unique fat and protein composition).***

***It is naturally rich in important nutrients such as Ca and vitamin A, and has a high bio-availability of iron.***

***It has a deliciously mild taste.***

***Kabrita contains our special DigestX OPO-fat blend (also known as beta-palmitate). Breast milk is known to contain a high amount of beta-palmitate and contributes to the typical breast-fed stools and calcium absorption.***

***Kabrita also contains prebiotic fibers: GOS / FOS. These fibers have a positive effect on the bacterial composition in the gut by stimulating growth of beneficial bacteria (Bifido- and lactobacilli).***

No evidence is given on the website to support any claims.

This milk meets current compositional requirements, and is more modified than other goats' milk based formula.

**Note:** The website for this brand of infant formula is based in Europe and the instructions for making up the milk on this website are not in line with current UK recommendations. However, the product sold in the UK does have appropriate labelling for making the product up safely, and these are the instructions that should be followed.

# Kendamil 1 First Infant Milk

This cows' milk based infant formula manufactured by Kendal Nutricare is currently available in the North West of England and in selected Morrisons and Asda supermarkets nationwide. The product is available in the UK as powder formulation only in 150g bag in box and 900g tins.

## Key components of this product

**TABLE 13**  
**Key components of Kendamil 1 First Infant Milk**

Kendamil 1 First Infant Milk	
Powdered infant milk (900g tin)	
<b>Composition</b>	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	Arachidonic acid, prebiotics, taurine, nucleotides
<b>Allergens</b>	Cows' milk,soya
<b>Protein source</b>	Cows' milk
<b>Whey:casein ratio</b>	60:40
<b>Fats used</b>	Sunflower, coconut, canola and algal oils (single cell oils) milk fat
<b>Suitability</b>	Suitable for vegetarians Not halal approved
<b>Cost per 100ml</b>	14p
<i>For comparison:</i> <b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on a powdered formulation)	10p

## Claims made

No specific health claims are made for this milk on the company website, however it does state that the product is made using whole milk.

Whilst whole milk is used in the product, after drying, it only accounts for about 16% of the dry weight of the powder, the majority being composed of whey powder, vegetable oils and lactose.

# Kendamil Mehadrin First Infant Milk

This cows' milk based infant formula manufactured by Kendal Nutricare has been specifically formulated and manufactured to comply with the beliefs and practices of Orthodox Jewish parents and carers. It is currently distributed via Global Kosher and is available only in stores they supply in London, Manchester and Gateshead. The infant milk is marketed as suitable for infants from birth to 12 months and is available as a powder formulation in 800g tins.

## Key components of this product

**TABLE 14**

**Key components of Kendamil Mehadrin First Infant Milk**

Kendamil Mehadrin First Infant Milk	
Powdered infant milk (800g tin)	
<b>Composition</b>	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	Arachidonic acid, taurine, nucleotides
<b>Allergens</b>	Cows' milk
<b>Protein source</b>	Cows' milk
<b>Whey:casein ratio</b>	60:40
<b>Fats used</b>	Rapeseed, coconut, sunflower and algal oils (single cell oils) milk fat
<b>Suitability</b>	Suitable for vegetarians Suitable for those following a Kosher diet. Not halal approved.
<b>Cost per 100ml</b>	19p
<i>For comparison:</i> <b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on a powdered formulation)	10p

## Claims made

No specific health claims are made for this milk on the company website, however it does state that the product is made using whole milk.

Whilst whole milk is used in the product, after drying, it only accounts for about 11% of the dry weight of the powder, the majority being composed of lactose, vegetable oils and skimmed milk powder.

# Mamia First Infant Milk

Mamia milk is marketed in ALDI stores and is sold as powder only in 900g tins.

## Key components of this product

TABLE 15

Key components of Mamia First Infant Milk

Mamia First Infant Milk	
Powdered infant milk 900g tin	
<b>Composition</b>	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	Arachidonic acid, prebiotics, taurine, nucleotides
<b>Allergens</b>	Cows' milk, soya, fish
<b>Protein source</b>	Cows' milk
<b>Whey:casein ratio</b>	60:40
<b>Fats used</b>	Palm, palm kernel, sunflower, rapeseed, fish and soya oils
<b>Suitability</b>	Not suitable for vegetarians Halal approved
<b>Cost per 100ml<sup>1</sup></b>	10p
<i>For comparison:</i>	10p
<b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on powdered formulation)	

There is currently no marketing information in the UK for this product but we will update this section if this becomes available.

The milk meets the compositional criteria required for infant milks.

# NANNYcare First Infant Milk

NANNYcare First Infant Milk is a goats' milk formula. It is available in powder form in 400g and 900g tins at some supermarkets and in specialist food shops.

## Key components of this product

**TABLE 16**  
**Key components of NANNYcare First Infant Milk**

NANNYcare First Infant Milk	
	Powdered infant milk
<b>Composition</b>	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	Taurine
<b>Allergens</b>	Goats' milk
<b>Protein source</b>	Goats' milk
<b>Whey:casein ratio</b>	20:80
<b>Fats used</b>	High oleic sunflower, rapeseed and sunflower oils
<b>Suitability</b>	Not suitable for vegetarians Halal approved
<b>Cost per 100ml</b>	33p
<i>For comparison:</i> <b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on a powdered formulation)	10p

## Claims made

### ***The only goat milk formula supported by clinical trials***

On the manufacturer's website, reference is made to a trial by Zhou et al (2014) which was used as evidence to EFSA (2012) when they reviewed the safety of goats' milk protein as a source of protein in infant formula. NANNYcare First Infant Milk was the formula used in that study. The EFSA recommendation was that milk from goats' milk or cows' milk can be a suitable protein source for infant formula provided the final product complies with the composition criteria laid down in the relevant EU Directive. The EFSA panel highlights that, if goat milk protein is used in infant formula, particular attention has to be given to the amino acid content by adding in appropriate free amino acids to ensure that the profile is adequate.

The following claims are also made:

**Goat milk:**

***Is an excellent source of high quality, readily digestible proteins.***

***Forms looser, softer and more porous curds in the baby's stomach. (This is particularly helpful to the baby's delicate developing digestive system.)***

***Has a casein profile which is closer to human milk (than cow's milk).***

***Has high levels of medium-chain fatty acids.***

***Has high levels of nucleotides.***

No evidence is provided on the website to support these claims.

In their evaluation of the suitability of goats' milk protein as the protein source in infant formula, EFSA clearly state that, while they note the differences in the composition of the caseins between goat and cow milk, no difference in digestibility has been observed (EFSA, 2012). In addition they state that there is no convincing evidence to support a lower incidence of allergic reactions in infants fed formula based on goats' milk protein compared with those fed cows' milk protein based formula.

The relevance of high levels of medium-chain fatty acids or nucleotides is not explained, but as stated previously, whether an infant formula is made from cows' or goats' milk protein, the composition has to meet current regulations.

## References

European Food Safety Authority (2012). *Scientific opinion on the suitability of goat milk protein as a source of protein in infant formulae and in follow-on formulae*. Parma, Italy: European Food Safety Authority. Available at: <http://www.efsa.europa.eu/en/efsajournal/pub/2603.htm>

Zhou SJ, Sullivan T, Gibson RA et al (2014). Nutritional adequacy of goat milk infant formula for term infants: a double blind randomised controlled trial. *British Journal of Nutrition* 111: 1641-51.

# Similac First Infant Milk

This cows' milk based infant formula manufactured by Abbott Nutrition is currently available in store and on-line from Boots Chemists only. The product is available as powder formulation in 850g tins.

## Key components of this product

**TABLE 17**

**Key components of Similac First Infant Milk**

Similac First Infant Milk	
Powdered infant milk (850g tin)	
<b>Composition</b>	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	Arachidonic acid, prebiotics, taurine, nucleotides
<b>Allergens</b>	Cows' milk, soya
<b>Protein source</b>	Cows' milk
<b>Whey:casein ratio</b>	48:52
<b>Fats used</b>	Sunflower, soya, coconut, fungal and algal oils (single cell oils)
<b>Suitability</b>	Suitable for vegetarians Halal approved
<b>Cost per 100ml</b>	16p
<i>For comparison:</i> <b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on a powdered formulation)	10p
<b>Preparation instructions</b>	Unlike other First Infant Milks this product is made up using one scoop of powder to 60ml of water. Other formula milks typically use one scoop of powder to 30ml of powder. The typical scoop weight for this product is 8.3g. This information is highlighted in red on the product packaging.

No specific health claims are made for this milk on the company website, however it does state that the product does not contain palm oil.

# SMA Pro First Infant Milk

SMA Pro First Infant Milk is available as powdered infant formula, a 70ml RTF (ready-to-feed) formulation and in 200ml and 1 litre RTF cartons.

## Key components of this product

**TABLE 18**  
**Key components of SMA Pro First Infant Milk**

SMA Pro First Infant Milk		
	Powder formulation	RTF (Ready to feed) 200 ml carton
<b>Composition</b>	Meets current regulations for infant formula composition	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	Prebiotics, taurine, arachidonic acid	Taurine, arachidonic acid, nucleotides
<b>Allergens</b>	Cows' milk, soya, fish	Cows' milk, soya, fish
<b>Protein source</b> <b>Whey:casein ratio</b>	Cows' milk 70:30	Cows' milk 65:35
<b>Fats used</b>	Palm, rapeseed, coconut and sunflower oils Fish and algal oils (single cell oils)	Palm, rapeseed, coconut and sunflower oils Fish and algal oils (single cell oils)
<b>Suitability</b>	Not suitable for vegetarians Halal approved	Not suitable for vegetarians Halal approved
<b>Cost per 100ml</b>	16p	40p
<i>For comparison:</i> <b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on a powdered formulation)	13p	13p

## What claims are made for SMA Pro?

**The quantity of protein has been reduced.**

SMA Nutrition claim to have used a “*new exclusive protein process*” to produce the protein component of their infant milk. No information is given on the website to explain what this process is, but in an email to First Steps Nutrition Trust, Nestlé Nutrition said that:

*“This patented process allows SMA Nutrition to reduce the total protein content in SMA PRO First Infant Milk to 1.25g/100ml (1.87g/100kcal) in line with scientific opinion while maintaining a desirable amino acid profile.”*

This infant milk claims to have the lowest protein content of any infant milk on the market. However, the difference is not sufficiently significant to differentiate it from all other brands; the difference in protein content between it and the brand with the next lowest protein content is 0.02g/100kcal (0.01g/100ml).

It has been suggested that a higher protein content in infant formula is associated with higher weight in the first two years of life (Koletzko et al, 2009). This association is thought to be a result of higher levels of circulating branched chain amino-acids in formula-fed infants compared to breastfed infants. Concentrations of these amino-acids are linked to insulin release, which in turn is linked to metabolic alterations which might be mechanisms for weight gain (Trabulsi et al, 2011). These two factors are being linked to suggest that lower protein milks may reduce weight gain in formula-fed infants, but whether protein plays a role in increased growth rate and higher BMI in childhood is still a matter of debate and requires considerably more research (EFSA, 2014).

Currently the four most commonly purchased infant formulas on the UK market all have around 1.3g protein/100ml. All infant formula must contain amino-acids in amounts specified in the regulations.

The SMA professional website claims that SMA Pro First Infant Milk is a:

***Nutritionally complete infant milk with a unique protein profile that is closer to breastmilk***

***SMA PRO First Infant Milk has lower levels of insulinogenic amino acids compared with other first infant milks.***

***Insulinogenic amino acids are shown to contribute to obesity later in life.***

There are, however, no articles from peer-reviewed journals to support these claims, and all of the claims above are referenced only as “*on file at Nestlé*”. We have asked for this data but it has not been provided.

Further claims made on the SMA Professional website include that SMA Pro First Infant Milk:

***Is the only First Infant Milk clinically proven to achieve a growth rate comparable with a breastfed baby.***

This claim is supported by one reference to a poster presentation by Grathwohl et al (2010) – all Nestlé employees – at a conference. This poster reported a meta-analysis of four studies looking at the growth of infants fed with Nan milk, but provides no data, and claims that there is good agreement with the WHO growth standard. However, this data does not appear to have been published subsequently in a peer-reviewed publication. It is unlikely that formula fed babies will have similar growth patterns to breastfed babies as it is

established internationally that infant formula-fed infants grow at a different pace to breastfed infants (Garza and de Onis, 2004).

There is no detail about which Nan formula the studies used, but it was one used in Australia, France and Italy before 2010. It is unclear whether SMA Pro First Infant Milk is the same milk as the one used in the studies which they say support this claim.

The website also claims that SMA Pro First Infant Milk is:

### ***Easy to digest***

This is supported by reference to the NHS Choices website which states that whey-based formula milks are suitable for newborns as they are thought to be easier to digest than casein-based formula milks (NHS Choices, 2014). This is not unique to SMA Pro as all infant milks based on cows' milk available in the UK are based on whey protein.

### ***Contains GOS/FOS for increased gastrointestinal comfort and softer stools similar to that of a breastfed infant.***

They provide references from two clinical trials to support this statement, one from Moro et al (2002) and the other from Vivatvakin et al (2010). In the study by Moro et al (2002), 90 healthy term infants were randomly allocated to receive formula milk supplemented with oligosaccharides at a concentration of 0.4g/100ml or 0.8g/100ml or placebo, over a period of 28 days. The infants receiving the formula milk supplemented with oligosaccharides showed a dose-dependent increase in the amount of *Bifidobacteria* in stools, but this may have no clinical significance. Stool softness was increased in the group receiving formula supplemented at 0.8g/100ml. The concentration of oligosaccharides which were used in the formula that resulted in greater stool softness was double that currently used in SMA Pro First Infant Milk.

Vivatvakin et al (2010) reported that infants fed a whey-based infant formula supplemented with oligosaccharides and long-chain polyunsaturated fatty acids had fewer hard stools and more soft stools than infants fed a casein-based formula with no additional ingredients. Infants fed the whey-based formula also had a microbiota composition and gastric and intestinal transit times closer to those of breastfed infants. The authors concluded that, based on parental reports of regurgitation, vomiting, crying and colic, whey-based formula supplemented with oligosaccharides and LCPs provides superior gastrointestinal comfort than casein-based formula that does not contain these added ingredients.

This trial was sponsored by Nestlé and does little to support the use of oligosaccharides in infant formula because the parental reported effects could not be attributed solely to the presence of GOS/FOS, as one of the trial formula milks was whey-based and the other casein-based. Current recommendations in the UK are that a first infant formula should be whey-based as whey is thought to be easier to digest than casein.

It is worth noting that only SMA Pro powdered formula contains GOS/FOS, not the ready-to-feed formula, suggesting that the company itself may not be convinced of the benefit of adding prebiotic: if it was, GOS/FOS would be added to all formula.

EFSA (2014) state that:

*“there is no evidence for health benefits from the addition of prebiotic oligosaccharides (GOS/FOS) to infant or follow-on formula.”*

***Nucleotides are no longer added to SMA Pro powder formulation.***

The powder formulation of SMA Pro First Infant Milk no longer contains nucleotides, but the RTF formula still contains them, suggesting the company is somewhat confused as to whether they think this is an ingredient to make claims about, or whether *not* putting nucleotides in is worth a claim as they are doing here.

EFSA (2014) are clear that:

*“there is no necessity to add nucleotides to infant or follow-on formula”.*

***Although LCPs can be made in the body, infants have a high demand for these nutrients. SMA PRO First Infant Milk contains Omega 3 and 6 LCPs.***

It is a regulatory requirement that infant formula provides a specific fatty acid profile, and it is not clear how the claim being made here about fatty acids in the milk relate to mandatory versus optional ingredients.

### **Our conclusion**

SMA Pro First Infant Milk meets the required compositional criteria for infant formula.

This infant formula offers no evidence-based advantage over any other first infant milk. The reduced protein emphasised on the website is only marginally lower than that found in other infant milks, and no evidence has been given to explain the new protein process which they say makes this formula ‘closer to breastmilk’.

The addition of prebiotics to the powdered version of SMA PRO means that this product is now more similar to other infant milks available on the market than the previous formulation, but the composition of the milks now vary by format (powder or RTF), which makes claims made confusing.

SMA Pro First Infant Milk is a mid-priced product, costing more than Cow & Gate and Hipp Organic Combiotic First Infant Milks, and less than Aptamil 1 First Milk or Aptamil Profutura 1 First Infant Milk. (For a list of the cost of various types of infant milks, see the Appendix.)

### **References**

EFSA (2014). *Scientific opinion on the composition of infant and follow-on formulae*. Parma, Italy: European Food Safety Authority. Available at <http://www.efsa.europa.eu/en/efsajournal/pub/3760>

Garza C, de Onis M for the WHO Multicentre Growth Reference Study Group (2004) Rationale for developing a new international growth reference. *Food and Nutrition Bulletin*, 25, (suppl 1) 55- 65.

Grathwohl D, Macé K, Fichot MC et al (2010). Evaluation of breastfed infants with respect to WHO growth standard: a meta-analysis. Poster presentation EAPS Congress. *Pediatric Research*; 68: 602.

Koletzko B, von Kries R, Closa R et al (2009). Lower protein in infant formula is associated with lower weight up to age 2 y: a randomized clinical trial. *American Journal of Clinical Nutrition*; 89: 1836-45.

Moro GE, Minoli I, Mosca M et al (2002). Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *Journal of Pediatric Gastroenterology and Nutrition*; 34: 291-95.

NHS Choices (2014). Types of infant formula. October 2014.  
<http://www.nhs.uk/Conditions/pregnancy-and-baby/Pages/types-of-infant-formula.aspx#close>. Accessed 28 September 2015.

Trabulsi J, Capeding R, Lebumfacil J et al (2011). Effect of an  $\alpha$ -lactalbumin-enriched infant formula with lower protein on growth. *European Journal of Clinical Nutrition*; 65 (2): 167-74.

Vivatvakin B, Mahayosnond A, Theambooniers A et al (2010). Effect of a whey-predominant starter formula containing LCPUFAs and oligosaccharides (FOS/GOS) on gastrointestinal comfort in infants. *Asia Pacific Journal of Clinical Nutrition*; 19(4): 473-480.

# SMA Wysoy

---

Soya protein based infant formula are permitted within the Infant Formula regulations, but in the UK the use of soya formula for infants under 6 months is not recommended without medical advice. There are a number of concerns about potential risks of soya formula use: soya proteins are potential allergens; the phyto-oestrogens present may have an impact on the future reproductive health of infants; the carbohydrate source is glucose syrup which is a greater risk for tooth decay; soya protein based formula have higher aluminium contents.

Only one brand of soya based formula is available in the UK. The product is not suitable for vegans as the vitamin D is sourced from sheep's wool lanolin. It is available in powdered form only in 860g tins.

## Key components of this product

**TABLE 19**  
**Key components of SMA Wysoy**

	<b>Powder formulation</b>
<b>Composition</b>	Meets current regulations for infant formula composition
<b>Unnecessary added ingredients</b>	Taurine, arachidonic acid
<b>Allergens</b>	Soya
<b>Protein source</b>	Soya
<b>Fats used</b>	Palm, soya, coconut and sunflower oils Algal oils (single cell oils)
<b>Carbohydrates used</b>	Glucose syrup
<b>Suitability</b>	Suitable for vegetarians Halal approved
<b>Cost per 100ml</b>	17p
<i>For comparison:</i> <b>Cheapest first infant milk currently available that meets all infant requirements/100ml</b> (based on a powdered formulation)	10p

No specific claims are made for Wysoy beyond the fact that it is lactose free and contains long chain fatty acids as required by the regulations.

## Hungry baby milks

---

There are four brands of hungry baby milks on the market: Aptamil Hungry Milk, Cow & Gate Infant Milk for Hungrier Babies, Hipp Organic Combiotic Hungry Infant Milk and SMA Extra Hungry. We have grouped these together as they are not recommended for use, and are all very similar.

Interestingly, as brands have altered their first infant milks, the hungry baby milks have not been changed, so either the companies do not consider these milks important enough to make the same changes to these as to their first infant milks, or they are not completely convinced by the evidence that they give for their own first milks. For example, despite SMA PRO and HiPP Organic Combiotic first milks making claims about the importance of lower protein in infant formula, they have not made changes to their hungry baby milks which both have 1.6g protein/100ml, an amount they claim is linked to higher weight gain. All hungry baby milks have a protein content considerably higher than first infant milks.

These hungry baby milks are casein-based and the manufacturers suggest that a whey:casein ratio of approximately 20:80 (which is similar to that in cows' milk) can result in slower gastric emptying, resulting in greater satiety. Evidence used to support this suggestion has, however, been from small studies of infants with reflux difficulties (Billeaud et al, 1990; Tolia et al, 1992) and these findings are not accepted as evidence of efficacy.

The manufacturers also suggest that the use of these milks may help delay weaning, but there is no scientific evidence to support this.

Cow & Gate suggest on their website that infants under 6 months may get a better night's sleep if they have hungry baby formula in the evening, but give no evidence to support this claim.

The nutritional composition of casein-dominant formulas is slightly different to that of whey-dominant infant formulas. However, the reported nutrient composition for hungry baby milks here is within recommended levels. A whey-based infant formula is recommended for use throughout the first year of life and therefore none of these milks should be recommended.

TABLE 20

## Summary of composition of hungry baby milks (powdered formulations only)

Nutrients per 100ml	Aptamil Hungry Milk	Cow & Gate Infant Milk for Hungrier Babies	Hipp Organic Combiotic Hungry Infant Milk	SMA Extra Hungry
Energy kcal	66	66	67	67
Protein g	1.6	1.6	1.6	1.6
Whey:casein ratio	20:80	20:80	20:80	20:80
Carbohydrate g	7.8	7.7	7.3	7.0
– of which lactose g	7.4	7.4	7.1	7.0
Source of added carbohydrate	Lactose, oligo-saccharides	Lactose, oligo-saccharides	Lactose, oligo-saccharides	Lactose
Fat g	3.1	3.1	3.5	3.6
Fat source	Palm, rapeseed, coconut and sunflower oils	Palm, rapeseed, coconut and sunflower oils	Palm, rapeseed and sunflower oils	Palm, coconut, sunflower and soya oils
Added LCPUFAs				
ARA	✓	✓	✓	✓
DHA	✓	✓	✓	✓
LCPUFA source	Fish and single cell oils	Fish and single cell oils	Fish and single sell oils	Single cell oils
Amounts of vitamins and minerals meet regulations	✓	✓	✓	✓
Prebiotics	✓	✓	✓	✗
Probiotics	✗	✗	✗	✗
Nucleotides	✓	✓	✗	✓
Taurine	✓	✓	✗	✓
L-carnitine	✓	✓	N/S	✗
Contains soya	✓	✓	✓	✓
Contains fish oil	✓	✓	✓	✗
Suitable for vegetarians	✗	✗	✗	✓
Halal approved	✓	✓	✗	✓
Cost per 100ml	17p	14p	14p	16p

**Note:** Check the packaging of RTF milks as these milks may vary in composition and suitability.

## References

Billeaud C, Guillet J, Sandler B (1990). Gastric emptying in infants with or without gastro-oesophageal reflux according to the type of milk. *European Journal of Clinical Nutrition*; 44: 577-83.

Tolia V, Lin C, Kuhns L (1992). Gastric emptying using three different formulas in infants with gastroesophageal reflux. *Journal of Pediatric Gastroenterology and Nutrition*; 15: 297-301.

## Appendix

### Cost of infant formula in the UK – July 2017

Costs of powdered infant formula have been calculated based on scoop weight and cost per 100ml prepared formula.

**TABLE 20. Cost of infant formula in the UK – July 2017**  
**Cost of starter packs of first infant formula**

Milk name	Weight	Cost (July 2017)	Where costed	Cost per 100ml made up milk	Cost per day birth – 2weeks 8 bottles per day (560ml)
Aptamil Profutura 1 First Milk RTF	6 x70 ml starter pack	£10.99	Boots	£2.61	£14.62
Aptamil 1 First Infant Milk	6 x 70ml starter pack	£8.00	Boots	£1.90	£10.64
SMA Pro First Milk RTF	6 x 70ml Starter pack	£7.00	Boots	£1.67	£9.35
Cow & Gate 1 First Milk RTF	6 x 70ml starter pack	£6.50	Boots	£1.55	£8.68

#### Cost of first infant milks: RTF formulations

Milk name	Weight	Cost (July 2017)	Where costed	Cost per 100ml made up milk	Cost per day based on average intake formula at 2-3 months (920ml/day)
Aptamil Profutura 1 First Infant Milk RTF	200ml	99p	Boots	50p	£4.60
Aptamil 1 First Infant Milk RTF	200ml	85p	Boots	43p	£3.96
Aptamil Hungry Milk RTF	200ml	85p	Boots	43p	£3.96
SMA Pro First Infant Milk RTF	200ml	80p	Boots	40p	£3.68
SMA Extra Hungry RTF	200ml	80p	Boots	40p	£3.68
HiPP Organic Combiotic First Infant Milk RTF	200ml	75p	Boots	38p	£3.50
HiPP Organic Combiotic Hungry Infant Milk RTF	200ml	75p	Ocado	38p	£3.50
Cow & Gate 1 First Infant Milk RTF	200ml	75p	Boots	38p	£3.50
Cow & Gate Infant Milk for hungrier babies RTF	200ml	75p	Boots	38p	£3.50
Aptamil 1 First Infant Milk RTF	1000ml	3.70	Boots	37p	£3.40

Milk name	Weight	Cost (July 2017)	Where costed	Cost per 100ml made up milk	Cost per day based on average intake formula at 2-3 months (920ml/day)
Aptamil Hungry Milk	1000ml	3.70	Boots	37p	£3.40
SMA Extra Hungry from birth RTF	1000ml	3.50	Boots	35p	£3.22
Cow & Gate 1 First Infant Milk RTF	1000ml	3.19	Boots	32p	£2.94
Cow & Gate Infant Milk for hungrier babies RTF	1000ml	3.19	Boots	32p	£2.94

### Cost of first infant milks: powdered formulations (PIF)

Milk name	Weight	Cost (July 2017)	Where costed	Cost per 100ml made up milk	Cost per day based on average intake formula at 2-3 months (920ml/day)
Holle Organic Infant Goat Milk Formula 1	400g	£10.95	Organic supplier	35p	£3.22
NANNYCare First Infant Milk (goats' milk based)	900g	£22.98	Organic supplier	33p	£3.04
Kabrita Gold 1 (goats' milk based)	800g	£19.95	Organic supplier	33p	£3.04
Holle Organic Infant Formula 1	400g	£8.85	Organic supplier	28p	£2.58
Aptamil Profutura 1 First Infant Milk	800g	£12.99	Boots	21p	£1.93
Kendamil Mehadrin First Infant Milk	800g	£13.99	Manufacturer Information	19p	£1.75
Aptamil 1 First Milk	900g	£11.00	Boots	17p	£1.56
Aptamil Hungry Milk	900g	£11.00	Boots	17p	£1.56
Similac First Infant Milk	850g	£10.95	Boots	17p	£1.47
SMA Wysoy	860g	£12.00	Boots	17p	£1.56
SMA Pro First Infant Milk	800g	£10.00	Boots	16p	£1.47
Castlemil First Milk	800g	£9.00	Manufacturer information	15p	£1.38
Cow & Gate 1 First Infant Milk	900g	£9.00	Boots	14p	£1.29
Cow & Gate Infant Milk for hungrier babies	900g	£9.00	Boots	14p	£1.29

<b>Milk name</b>	<b>Weight</b>	<b>Cost (July 2017)</b>	<b>Where costed</b>	<b>Cost per 100ml made up milk</b>	<b>Cost per day based on average intake formula at 2-3 months (920ml/day)</b>
HiPP Organic Combiotic Hungry Infant Milk	800g	£8.50	Boots	14p	£1.29
Kendamil 1 First Milk	900g	£9.89	Manufacturer information	14p	£1.29
HiPP Organic Combiotic First Infant Milk	800g	£8.50	Boots	13p	£1.20
Bebivita First Infant Milk	500g	£3.99	LIDL (only available at this supermarket)	10p	92p
Mamia First Infant Milk	900g	£6.99	ALDI (only available at this supermarket)	10p	92p

